

132	HFTBM38	546	Inhibition of squalene synthetase gene transcription.	Reporter Assay: construct contains regulatory and coding sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the	158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
-----	---------	-----	---	---	--	---

				<p>contents of which are herein incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72 hours. HepG2 is a human hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.</p>	
132	HFTBM38	546	CD152 in Human T cells	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis,</p>
132	HFTBM38	546	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>		

				<p>activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative</p>
--	--	--	--	---	--

				disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
133	HIFVGK35	547	Production of IL-6	IL-6 F/MAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal
				A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention

				<p>immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and</p>	<p>includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include</p>
--	--	--	--	--	--

				<p>differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes</p>
--	--	--	--	---	--

133	HFVGK35	547	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to</p>	<p>mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications</p>
-----	---------	-----	---------------------	---	--

				<p>test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma,</p>
--	--	--	--	---	--

					prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
134	HFXBN86	548	Production of MIP1alpha	<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,</p>

				<p>cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and</p>	<p>multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma,</p>
--	--	--	--	---	---

					upregulate T cell proliferation and functional activities.	prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
135	HFXBT66	549	IL-2 in Human T cells			
135	HFXBT66	549	IFNg in Human T-cell 293T			
136	HFXFZ46	550	HLA-DR in Human T cells			
137	HGBER72	551	Activation of transcription through AP1 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as	

			<p>and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is an IL-2 dependent suspension culture cell line that also responds to IL-4.</p>	<p>described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis,</p>
--	--	--	---	--

137	HGBER72	551	<p>Activation of transcription through NFAT response in immune cells (such as T-cells).</p> <p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA</p>	<p>suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p> <p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as</p>
-----	---------	-----	--	---

				<p>85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---	---

137	HGBER/72	551	<p>Activation of transcription through NFKB response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic,</p>
-----	----------	-----	---	--	--

				in its entirety. Exemplary human T cells, such as the MOLT4, that may be used according to these assays are publicly available (e.g., through the ATCC).	esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.
137	HGBER72	551	SEAP in Senescence Assay		
138	HGBEY14	552	Production of TNF alpha by dendritic cells	TNFα FMA T. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative

			<p>other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890</p>	<p>highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>
--	--	--	--	--

				<p>(1198); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>“Hyperproliferative Disorders”). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and</p>
--	--	--	--	--	---

					asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
138	HGBEY14	552	IFNg in Human T-cell 2B9		
139	HGBGN34	553	SEAP in HIB/CRE		
139	HGBGN34	553	Production of IL-5	IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis.

			<p>evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human</p>	<p>An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative</p>
--	--	--	---	--

				lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
139	HGBGN34	553	SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)		
139	HGBGN34	553	Activation of Transcription	Assays for activation of transcription are well-known in the art and may be used and routinely modified to assess ability of polypeptides of the invention to inhibit or activate transcription. An example of such an assay follows: Cells were pretreated with SID	

140	HGLBG15	554		<p>supernatants or controls for 15-18 hours. SEAP activity was measured after 48 hours.</p> <p>LS174T is an epithelial colon adenocarcinoma cell line. Its tumorigenicity in nude mice make cell line LS174T a model for studies on the mechanism of synthesis and secretion of specific tumoral markers in colon cancer. See, Patan et al., Circ Res, 89(8):732-39 (2001), the contents of which are herein incorporated by reference in its entirety.</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-</p>
			<p>Production of IL-10 and activation of T-cells.</p>	<p>Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention)</p>	

				<p>to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p>	mediated immune response.
--	--	--	--	---	---------------------------

141	HHEGS55	555	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred</p>
-----	---------	-----	--	---	---

				<p>available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of</p>
--	--	--	--	---	--

					immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
142	HHEOW19	556	Regulation of viability or proliferation of immune cells (such as human eosinophil EOL-1 cells).	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of eosinophil cells and cell lines. For example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells.	Highly preferred indications include eosinophilia, asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation.

				<p>Eosinophils are a type of immune cell important in allergic responses; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Eosinophil cell lines that may be used according to these assays are publicly available and/or may be routinely generated. Exemplary eosinophil cells that may be used according to these assays include EOL-1 Cells.</p>	<p>Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>
142	HHEOW19	556	Production of MIP1alpha	<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p>

			<p>macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells</p>	<p>Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>
--	--	--	---	---

142	HHEOW19	556		<p>are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>“Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>Production of TNF alpha by dendritic cells</p>	<p>TNFα FMT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under “Immune Activity”, “Blood-Related Disorders”, and/or “Cardiovascular Disorders”), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,</p>

				<p>evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells</p>	<p>Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	---	--

				are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
142	HHEOW19	556	Insulin Secretion	Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy

			secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly	and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract
--	--	--	---	---

				<p>available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
143	HHFEC39	557	<p>Production of IL-10 and activation of T-cells.</p>	<p>Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells.</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease,</p>

				<p>Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary</p>	<p>multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	---	---

144	HHFF87	558	<p>Regulation of viability or proliferation of immune cells (such as human eosinophil EOL-1 cells).</p>	<p>T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p> <p>Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of eosinophil cells and cell lines. For example, the CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Eosinophils are a type of immune cell important in allergic responses; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction.</p>	<p>Highly preferred indications include eosinophilia, asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications</p>
-----	--------	-----	---	---	---

				Eosinophil cell lines that may be used according to these assays are publicly available and/or may be routinely generated. Exemplary eosinophil cells that may be used according to these assays include EOL-1 Cells.	include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.
144	HHFFF87	558	CD71 in Human T cells		
145	HHFFL34	559	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm,	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below),

				<p>boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia,</p>
			<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	

					pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
145	HHFFL34	559	SEAP in HIB/CRE		
145	HHFFL34	559	IL-10 in Human T-cell 293T		
146	HHFFS40	560	Endothelial Cell Apoptosis	Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for

				<p>agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an</p>	<p>inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly</p>
--	--	--	--	--	---

				<p>example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>preferred indications include neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly</p>
--	--	--	--	---	---

					<p>preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer.</p> <p>Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery</p>
--	--	--	--	--	--

					<p>disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions.</p>
--	--	--	--	--	--

					Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
146	HHFFS40	560	Production of TNF alpha by dendritic cells	TNFa FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method

				<p>cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1198); Dahlen et al., J Immunol 160(7):3585-3593 (1998);</p>	<p>for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred</p>
--	--	--	--	---	---

				<p>Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is</p>
--	--	--	--	--	---

146	HHFFS40	560	Stimulation of Calcium Flux in pancreatic beta cells.	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al.,	infection (e.g., an infectious disease as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below),
-----	---------	-----	---	---	--

				<p>Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 (Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	--

147	HHGCS78	561	<p>Regulation of transcription via DMEF1 response element in adipocytes and pre-adipocytes</p>	<p>Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays</p>	<p>A highly preferred indication is diabetes mellitus. Additional highly preferred indications include complications associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision</p>
-----	---------	-----	--	--	---

			disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to	impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
--	--	--	--	--

				adipose-like conversion under appropriate differentiation culture conditions.	
147	HHGCS78	561	Glucose Production in H4IIE		
147	HHGCS78	561	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke

148	HHGDT26	562	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>endothelial cells (MVEC).</p> <p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma,</p>
-----	---------	-----	--	---	---

				<p>contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
148	HHGDT26	562	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production.</p>

				<p>invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for</p>
--	--	--	--	---	---

				<p>killer cell line with cytolytic and cytotoxic activity.</p> <p>example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious</p>
--	--	--	--	--

149	HHPFU28	563	Proliferation of immune cells (such as the HMC-1 human mast cell line)	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of eosinophil cells and cell lines. For example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Mast cells are found in connective and mucosal tissues throughout the body. Mast cell activation (via immunoglobulin E -antigen, promoted by T helper cell type 2 cytokines) is an important component of allergic disease. Dysregulation of mast cell apoptosis may play	disease as described below under "Infectious Disease"). Highly preferred indications include asthma, allergy, mastocytosis (a rare, heterogeneous disorder characterized by excessive accumulation of mast cells, and their proliferation and action in the skin, central nervous system, and other organs). Preferred indications also include hematopoietic and immunological disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), infection (e.g., as described below under "Infectious Disease"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below).
-----	---------	-----	--	---	--

				<p>a role in allergic disease and mast cell tumor survival. Mast cell lines that may be used according to these assays are publicly available and/or may be routinely generated. Exemplary mast cells that may be used according to these assays include HMC-1, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	
149	HHPFU28	563	Myoblast cell proliferation	<p>Assays for muscle cell proliferation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit myoblast cell proliferation. Exemplary assays for myoblast cell proliferation that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays</p> <p>Highly preferred indications include diabetes, myopathy, muscle cell atrophy, cancers of muscle (such as, rhabdomyoma, and rhabdosarcoma), cardiovascular disorders (such as congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, vascular disease, and also as described below under "Cardiovascular Disorders"), stimulating myoblast proliferation, and inhibiting myoblast proliferation.</p>	

				disclosed in: Soeta, C., et al. "Possible role for the c-ski gene in the proliferation of myogenic cells in regenerating skeletal muscles of rats" Dev Growth Differ Apr;43(2):155-64 (2001); Ewton DZ, et al., "IGF binding proteins-4, -5 and -6 may play specialized roles during L6 myoblast proliferation and differentiation" J Endocrinol Mar;144(3):539-53 (1995); and, Pampusch MS, et al., "Effect of transforming growth factor beta on proliferation of L6 and embryonic porcine myogenic cells" J Cell Physiol Jun;143(3):524-8 (1990); the contents of each of which are herein incorporated by reference in their entirety. Exemplary myoblast cells that may be used according to these assays include the rat myoblast L6 cell line. Rat myoblast L6 cells are an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuse to form multinucleated myotubes and striated fibers after culture in differentiation media.	
--	--	--	--	--	--

149	HHPFU28	563	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred</p>
-----	---------	-----	--	---	---

				<p>available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of</p>
--	--	--	--	---	--

					immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
149	HHPFU28	563	SEAP in HIB/CRE		
149	HHPFU28	563	Activation of Endothelial Cell ERK Signaling Pathway.	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for</p>	

			agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Berra et al., Biochem Pharmacol 60(8):1171-1178 (2000); Gupta et al., Exp Cell Res 247(2):495-504 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.	stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell differentiation. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for
--	--	--	---	--

					<p>reducing cardiac hypertrophy.</p> <p>An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”).</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the</p>
--	--	--	--	--	---

				<p>arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	---

				hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph

				known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Rolfe BE, et al., Atherosclerosis, 149(1):99-110 (2000); Panettieri RA Jr, et al., J Immunol, 154(5):2358-2365 (1995); and, Grunstein MM, et al., Am J Physiol Lung Cell Mol Physiol, 278(6):L1154-L1163 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include Aortic Smooth Muscle Cells (AOSMC); such as bovine AOSMC.	polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Vascular Disease, Atherosclerosis, Restenosis, Stroke, and Asthma.
150	HHsBI65	564	Regulation of apoptosis in	Caspase Apoptosis. Assays for caspase apoptosis are well	A highly preferred indication is diabetes mellitus.

			pancreatic beta cells.	<p>known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Apoptosis in pancreatic beta is associated with induction and progression of diabetes. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Loweth, AC, et al., FEBS Lett, 400(3):285-8 (1997); Saini, KS, et al., Biochem Mol Biol Int, 39(6):1229-36 (1996); Krautheim, A., et al., Br J Pharmacol, 129(4):687-94 (2000); Chandra J, et al., Diabetes, 50 Suppl 1:S44-7 (2001); Suk K, et al., J Immunol, 166(7):4481-9 (2001); Tejedo J, et al., FEBS Lett, 459(2):238-43 (1999); Zhang, S., et al., FEBS Lett,</p>	<p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness),</p>
--	--	--	------------------------	---	--

				<p>455(3):315-20 (1999); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated.</p> <p>Exemplary pancreatic cells that may be used according to these assays include RIN-m. RIN-m is a rat adherent pancreatic beta cell insulinoma cell line derived from a radiation induced transplantable rat islet cell tumor. The cells produce and secrete islet polypeptide hormones, and produce insulin, somatostatin, and possibly glucagon. ATTC: #CRL-2057</p> <p>Chick et al. Proc. Natl. Acad. Sci. 1977 74:628; AF et al. Proc. Natl. Acad. Sci. 1980 77:3519.</p>	<p>ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
151	HHSDI53	565	Activation of transcription through	Assays for the activation of transcription through the Serum	A preferred embodiment of the invention includes a method

			<p>serum response element in immune cells (such as T-cells).</p>	<p>Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T</p>	<p>for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include</p>
--	--	--	--	--	--

				<p>cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues,</p>
--	--	--	--	---	--

151	HHSDI53	565	Production of IL-5	<p>IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins</p>	<p>hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis.</p>
-----	---------	-----	--------------------	---	---

			<p>evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human</p>	<p>An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative</p>
--	--	--	---	--

				lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
151	HHSD153	565	SEAP in SW480		
152	HHSFC09	566	Production of IFNgamma using a T cells	IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for	A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood

				<p>immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al.,</p>	<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative</p>
--	--	--	--	--	--

			<p>"Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---

153	HHSGL28	567	HLA-DR in Human T cells	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.
154	HISBA38	568	Production of IL-10 and activation of T-cells.		

				<p>incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p>	
155	HJMAA03	569	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related</p>

			<p>for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast,</p>
--	--	--	--	---

	HJMAA03	569	Stimulation of	Assays for measuring secretion	<p>lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred</p>
--	---------	-----	----------------	--------------------------------	---

155	insulin secretion from pancreatic beta cells.	<p>of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-</p>	<p>indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic</p>
-----	---	--	---

			<p>204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
156	HJMAV41	570	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as</p>

				<p>the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g.,</p>
--	--	--	--	---	--

					<p>malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	--	--

156	HJMAV41	570	TNF α in Human T cells		
157	HJMAV90	571	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and

				<p>by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease,</p>
--	--	--	--	---	---

157	HJMA Y90	571	Production of IL-6	<p>IL-6 FMA T. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by</p>	<p>inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described</p>
-----	----------	-----	--------------------	--	---

				<p>cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925</p>	<p>below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach,</p>
--	--	--	--	--	--

				<p>(1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
157	HJMA Y90	571	<p>Production of VCAM in endothelial cells (such as human</p>	<p>Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to</p>	<p>Highly preferred indications include inflammation (acute and chronic), restenosis, atherosclerosis, asthma and</p>

			umbilical vein endothelial cells (HUVEC))	<p>assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory</p>	<p>allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	---	--	---

159	HJPBK28	573	<p>Activation of transcription through NFKB response element in immune cells (such as T-cells).</p>	<p>responses.</p> <p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia,</p>
-----	---------	-----	---	---	--

				<p>contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
160	HJPC08	574	<p>Activation of T-Cell p38 or JNK Signaling Pathway.</p>	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as</p>

				<p>art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the</p>	<p>described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	--	---

				ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
161	HKABU43	575	Activation of Adipocyte ERK Signaling Pathway	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely	A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte

				<p>modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like</p>	<p>differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described</p>
--	--	--	--	---	--

				conversion under appropriate differentiation conditions known in the art.	below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular
--	--	--	--	---	--

					Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis,
--	--	--	--	--	--

				degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
161	HKABU43	575	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to
				Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below),

				<p>assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2</p>	<p>immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	--	--

					polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
162	HKACI79	576		SEAP in HIB/CRE		
162	HKACI79	576		CD152 in Human T cells		
162	HKACI79	576		HLA-DR in Human T cells		
162	HKACI79	576		IFNg in Human T-cell 2B9		
162	HKACI79	576		Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or

			<p>antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia,</p>
--	--	--	--	--

					acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
163	HKAFF50	577	CD69 in Human T cells	Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element	A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment
163	HKAFF50	577	Activation of transcription through CD28 response element in immune cells (such as T-cells).		

				<p>activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. An additional highly preferred indication includes infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms</p>
--	--	--	--	--	---

					and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication is infection (e.g., tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune
--	--	--	--	--	--

					<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
164	HKGBF25	578	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>

				<p>SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>"Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic,</p>
--	--	--	--	--	---

					<p>esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
	HKGBF25	578	SEAP in HepG2/Squale-		

			synthetase(stimulati on)			
165	HKMLK03	579	Inhibition of squalene synthetase gene transcription.	Reporter Assay: construct contains regulatory and coding sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the contents of which are herein incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72 hours. HepG2 is a human hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.		A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of
165	HKMLK03	579	CD69 in Human T cells			
166	HKMLM95	580	Production of IFNgamma using a T cells	IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage		

				<p>activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J</p>	<p>IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma,</p>
--	--	--	--	--	---

				<p>Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	---	---

167	HLDBG17	581	Activation of T-Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol</p>	<p>asthma and allergy.</p> <p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma,</p>
-----	---------	-----	--	---	--

				<p>Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
168	HLDCA54	582	RANTES in Human T cells		
169	HLDQU79	583	Regulation of viability and proliferation of pancreatic beta cells.	Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease

				<p>and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely</p>	<p>(e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below,</p>
--	--	--	--	---	--

				<p>generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
169	HLDQU79	583	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis,</p>

			<p>antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative</p>
--	--	--	--	--

					disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
170	HLDRT09	584	SEAP in Alk Phos C2C12		
170	HLDRT09	584	Activation of transcription through serum response element in immune	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An

			<p>cells (such as T-cells).</p>	<p>be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the</p>	<p>alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as</p>
--	--	--	---------------------------------	--	--

				<p>CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>described below under “Hyperproliferative Disorders”). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
--	--	--	--	--	--

					meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
170	HLDRT09	584	IgG in Human B cells SAC		
171	HLHAP05	585	Production of MIP1alpha	<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,</p>

			<p>cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and</p>	<p>multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma,</p>
--	--	--	---	---

				upregulate T cell proliferation and functional activities.	prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
171	HLHAP05	585	Regulation of transcription through the FAS promoter element in hepatocytes	Assays for the regulation of transcription through the FAS promoter element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the FAS promoter element in a reporter construct and to regulate transcription of FAS, a key enzyme for lipogenesis. FAS promoter is regulated by many transcription factors including SREBP. Insulin increases FAS gene transcription in livers of diabetic mice. This stimulation of transcription is also somewhat glucose dependent. Exemplary assays that may be used or routinely modified to	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,</p>

				<p>test for FAS promoter element activity (in hepatocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Xiong, S., et al., Proc Natl Acad Sci U.S.A., 97(8):3948-53 (2000); Roder, K., et al., Eur J Biochem, 260(3):743-51 (1999); Oskouian B, et al., Biochem J, 317 (Pt 1):257-65 (1996); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays, such as H4IIE cells, are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays include rat liver hepatoma cell line(s) inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
HLHCS23	586	Regulation of	Assays for the regulation of	A highly preferred indication is	

172	viability and proliferation of pancreatic beta cells.	<p>viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ohtani KI, et al., Endocrinology, 139(1):172-8 (1998); Krautheim A, et al, Exp Clin Endocrinol Diabetes, 107 (1):29-34 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that</p>	<p>diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic</p>
-----	---	---	---

				<p>may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
172	HLHCS23	586	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include</p>

			<p>response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma,</p>
--	--	--	---	--

					<p>melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below</p>
--	--	--	--	--	---

172	HLHCS23	586	Production of IFNgamma using a T cells	<p>IFNgamma FMA.T. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon</p>	<p>under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity"; "Blood-Related Disorders"; and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T</p>
-----	---------	-----	--	---	---

			<p>gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or</p>	<p>cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
--	--	--	---	---

				CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
172	HLHCS23	586	VEGF in SW480		
173	HLIBO72	587	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,

			antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.	<p>Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	---

					<p>conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
174	HLICE88	588	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for</p>

				<p>(including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred</p>
--	--	--	--	---	--

					<p>indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An</p>
--	--	--	--	--	---

174	HLICE88	588	Stimulation of insulin secretion from pancreatic beta cells.	Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et	additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular
-----	---------	-----	--	---	--

			<p>al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety.</p> <p>Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated.</p> <p>Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
175	HLICO10	589	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to</p> <p>Activation of transcription through serum response element in immune cells (such as T-</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred</p>

			cells).	<p>assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2</p>	<p>embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>
--	--	--	---------	--	--

				<p>dependent suspension culture of T cells with cytotoxic activity.</p>	<p>“Hyperproliferative Disorders”). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	---	--

					cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
175	HLICO10	589	Activation of Transcription	Assays for activation of transcription are well-known in the art and may be used and routinely modified to assess ability of polypeptides of the invention to inhibit or activate transcription. An example of such an assay follows: Cells were pretreated with SID supernatants or controls for 15-18 hours. SEAP activity was measured after 48 hours. LS174T is an epithelial colon adenocarcinoma cell line. Its tumorigenicity in nude mice make cell line LS174T a model for studies on the mechanism of synthesis and secretion of specific tumoral markers in colon cancer. See, Patan et al., Circ Res, 89(8):732-39 (2001), the contents of which are herein incorporated by reference in its entirety.	
176	HLJBS28	590	Activation of transcription through	Assays for the activation of transcription through the Serum	A preferred embodiment of the invention includes a method

			serum response element in immune cells (such as T-cells).	<p>Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T</p>	<p>for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include</p>
--	--	--	---	--	--

				<p>cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues,</p>
--	--	--	--	---	--

					hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
177	HLMBW89	591	Upregulation of CD154 and activation of T cells	<p>CD154 FMAT. CD154 (a.k.a., CD40L) expression is induced following activation of T cells. Interaction between CD154 and CD40 on B cells is required for correct antibody class switching and germinal center formation. Mutations in CD154 are linked to immunodeficiencies and increased susceptibility to infections. Assays for immunomodulatory proteins important for antibody class switching and TH1 function and expressed on activated T helper lymphocytes are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, modulate</p>	<p>A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g.,</p>

				<p>antibody class switching, mediate TH1 function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD154, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Mackey et al., J Leukoc Biol 63(4):418:428 (1998); and Skov et al., 164(7):3500-3505 (2000), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human</p>	<p>AIDS). Preferred indications include boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune response. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune</p>
--	--	--	--	---	---

178	HLMGP50	592	HLA-DR in Human T cells	<p>lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p> <p>HLA-DR FMAT. MHC class II is essential for correct presentation of antigen to CD4+ T cells. Deregulation of MHC class II has been associated with autoimmune diseases (e.g., diabetes, rheumatoid arthritis, systemic lupus erythematosus, and multiple sclerosis). Assays for immunomodulatory proteins expressed on MHC class II expressing T cells and antigen presenting cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of MHC class II products, such as</p>	<p>reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, inflammation and inflammatory disorders, and asthma and allergy.</p>
				<p>Immune</p> <p>Highly preferred indications include immunological disorders such as described herein under the heading "Immune Activity" and/or "Blood-Related Disorders" (particularly including, but not limited to, immune disorders involving T-cells). Highly preferred embodiments of the invention include methods of preventing, detecting, diagnosing, treating and/or ameliorating disorders of the immune system (particularly including, but not limited to, immune disorders involving T-cells).</p>	

				<p>HLA-DR antigens, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Lamour et al., Clin Exp Immunol 89(2):217-222 (1992); Hurme and Silvola, Immunol Lett 20(3):217-222 (1989); Gansbacher and Zier, Cell Immunol 117(1):22-34 (1988); and Itoh et al., J Histochem Cytochem 40(11):1675-1683, the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate</p>
--	--	--	--	---

179	HLMJB64	593	Activation of transcription through GAS response element in immune cells (such as eosinophils).	humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors. Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate gene expression (commonly via STAT transcription factors) involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al.,	Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting an eosinophil-mediated immune response and, alternatively, suppressing an eosinophil-mediated immune response.
-----	---------	-----	---	--	---

				<p>Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995); the contents of each of which are herein incorporated by reference in its entirety. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate or inhibit activation of immune cells include assays disclosed and/or cited in: Mayumi M., "EoL-1, a human eosinophilic cell line" Leuk Lymphoma; Jun;7(3):243-50 (1992); Bhattacharya S, "Granulocyte macrophage colony-stimulating factor and interleukin-5 activate STAT5 and induce CIS1 mRNA in human peripheral blood eosinophils" Am J Respir Cell Mol Biol; Mar;24(3):312-6 (2001); and, Du J, et al., "Engagement of the CrkL adapter in interleukin-5 signaling in eosinophils" J Biol</p>
--	--	--	--	---

				<p>Chem; Oct 20;275(42):33167-75 (2000); the contents of each of which are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are a type of immune cell important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Increases in GAS mediated transcription in eosinophils is typically a result of STAT activation, normally a direct consequence of interleukin or other cytokine receptor stimulation (e.g. IL3, IL5 or GMCSF).</p>	
179	HLMJB64	593	Activation of transcription through NFkB response element in immune cells (such as mast cells).	<p>This reporter assay measures activation of the NFkB signaling pathway in HMC-1 human mast cell line. Activation of NFkB in mast cells has been linked to production of certain cytokines, such as IL-6 and IL-9. Assays for the activation of transcription through the NFkB response element are well-known in the art and may be</p> <p>Highly preferred indication includes allergy, asthma, and rhinitis. Additional highly preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include immunological and hematopoietic disorders (e.g.,</p>	

				<p>used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFκB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFκB response element that may be used or routinely modified to test NFκB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Stassen et al, J Immunol 166(7):4391-8 (2001); and Marquardt and Walker, J Allergy Clin Immunol 105(3):500-5 (2000), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available</p>	<p>as described below under "Immune Activity", and "Blood-Related Disorders"). Preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancer, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary tract cancers and as described below under "Hyperproliferative Disorders".</p>
--	--	--	--	---	---

				(e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.	
180	HLQAS12	594	Production of GM-CSF	GM-CSF FMAT. GM-CSF is expressed by activated T cells, macrophages, endothelial cells, and fibroblasts. GM-CSF regulates differentiation and proliferation of granulocytes-macrophage progenitors and enhances antimicrobial activity in neutrophils, monocytes and macrophage. Additionally, GM-CSF plays an important role in the differentiation of dendritic cells and monocytes, and increases antigen presentation. GM-CSF is considered to be a proinflammatory cytokine. Assays for immunomodulatory proteins that promote the production of GM-CSF are well known in the art and may be used or routinely modified to	A highly preferred embodiment of the invention includes a method for stimulating the production of GM-CSF. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of GM-CSF. Highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., as described below under "Infectious Disease". Highly preferred indications include blood disorders (e.g., neutropenia (and the prevention of neutropenia (e.g., in HIV infected patients), and/or as described below under "Immune

			<p>assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and modulate the growth and differentiation of leukocytes. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as GM-CSF, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Ye et al., J Leukoc Biol (58(2):225-233, the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include asthma. Highly preferred indications include neoplastic diseases (e.g., leukemia (e.g., acute lymphoblastic leukemia, and acute myelogenous leukemia), lymphoma (e.g., non-Hodgkin's lymphoma and Hodgkin's disease), and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	--

				<p>ATCC) or may be isolated using techniques disclosed herein or otherwise known in the art. Natural killer (NK) cells are large granular lymphocytes that have cytotoxic activity but do bind antigen. NK cells show antibody-independent killing of tumor cells and also recognize antibody bound on target cells, via NK Fc receptors, leading to cell-mediated cytotoxicity.</p>	<p>hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications include: suppression of immune reactions to transplanted organs and tissues (e.g., bone marrow transplant); accelerating myeloid recovery; and mobilizing hematopoietic progenitor cells. Preferred indications include boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune response. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and allergy.</p>
181	HLQCL64	595	CD69 in Human T cells		
182	HLWAV47	596	Activation of transcription through	Assays for the activation of transcription through the Serum	A preferred embodiment of the invention includes a method

			<p>serum response element in immune cells (such as T-cells).</p>	<p>Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T</p>	<p>for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include</p>
--	--	--	--	--	--

				<p>cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues,</p>
--	--	--	--	---	--

183	HLWBB73	597	Production of IL-6	<p>IL-6 FMT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention</p>	<p>hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus</p>
-----	---------	-----	--------------------	---	---

				<p>(including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these</p>	<p>erythematosis, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	---	--

				assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
184	HLWCN37	598	Production of IFNgamma using a T cells	IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC	A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred

			<p>expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-</p>	<p>indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described</p>
--	--	--	--	--

			<p>204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---

185	HL YEU59	599	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach,</p>
-----	----------	-----	--	---	---

				<p>in its entirety. Mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is an IL-2 dependent suspension culture cell line that also responds to IL-4.</p>	<p>brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
186	HLYGB19	600	CD152 in Human T cells	<p>CD152 FMAT. CD152 (a.k.a. CTLA-4) expression is restricted to activated T cells. CD152 is a negative regulator of T cell proliferation. Reduced CD152 expression has been linked to hyperproliferative and autoimmune diseases. Overexpression of CD152 may lead to impaired</p>	<p>Immune</p> <p>Highly preferred indications include immunological disorders such as described herein under the heading "Immune Activity" and/or "Blood-Related Disorders" (particularly including, but not limited to, immune disorders involving T-cells). Highly preferred embodiments of the invention include methods of preventing, detecting, diagnosing,</p>

				<p>immunoresponses. Assays for immunomodulatory proteins important in the maintenance of T cell homeostasis and expressed almost exclusively on CD4+ and CD8+ T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate the activation of T cells, maintain T cell homeostasis, and/or mediate humoral or cell-mediated immunity.</p> <p>Exemplary assays that test for immunomodulatory proteins evaluate the upregulation of cell surface markers, such as CD152, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include, for example, the assays disclosed in Miraglia et al., J</p>	<p>treating and/or ameliorating disorders of the immune system (particularly including, but not limited to, immune disorders involving T-cells).</p>
--	--	--	--	--	--

				<p>Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); McCoy et al., Immunol Cell Biol 77(1):1-10 (1999); Oostervegal et al., Curr Opin Immunol 11(3):294-300 (1999); and Saito T, Curr Opin Immunol 10(3):313-321 (1998), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>A highly preferred embodiment of the invention includes a method for increasing</p>
186	HLYGB19	600	Activation of Skeletal Muscle Cell PI3 Kinase	<p>Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal</p>	

			<p>Signalling Pathway</p>	<p>transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for P13 kinase activity that may be used or routinely modified to test P13 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays</p>	<p>muscle cell survival. An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is inhibited. Highly preferred indications include disorders of the musculoskeletal system.</p>
--	--	--	---------------------------	--	---

				include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage
--	--	--	--	--	--

					<p>(e.g. due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hypoglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications</p>
--	--	--	--	--	---

					<p>associated with obesity.</p> <p>Additional highly preferred indications include weight loss or alternatively, weight gain.</p> <p>Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal system including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, and vascular disease.</p> <p>Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	--	--

					conditions, such as, hyperplasia, metaplasia, and/or dysplasia.
187	HL YGE16	601	SEAP in Molt4/SRE		
187	HL YGE16	601	SEAP in NK16/STAT6		
188	HL YGY91	602	Insulin Secretion	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al.,</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other</p>

				<p>Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219:</p>	<p>diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---	--

188	HLYG91	602	Production of IL-13 and activation of T-cells.	<p>547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Assays for production of IL-13 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-13 and/or activation of T-cells.</p> <p>Exemplary assays for IL-13 production that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays such as disclosed and/or cited in: Grunig, G, et al., "Requirement for IL-13 independently of IL-4 in Experimental asthma" Science; 282: 2261-2263 (1998), and Wills-Karp M, et al., "Interleukin-13: central mediator of allergic asthma" Science; 282: 2258-2261</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
-----	--------	-----	--	--	--

				(1998); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL13, a Th2 type cytokine, is a potent stimulus for mucus production, airway hyper-responsiveness and allergic asthma. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
189	HMCFH60	603	Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions.	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious

			<p>Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia,</p>
--	--	--	--	---

					thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
189	HMC6H60	603	SEAP in Jurkat/IL4 promoter		
189	HMC6H60	603	SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)		
189	HMC6H60	603	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications

				<p>the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other</p>
--	--	--	--	--	---

					<p>preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
189	HMC6H60	603	<p>Activation of transcription through GAS response element in immune</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>

			<p>well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary</p>	<p>“Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt’s lymphoma, non-Hodgkins lymphoma, Hodgkin’s disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as</p>
--	--	--	--	---

				human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).	described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
	HMCFH60	603	Activation of	Assays for the activation of	A highly preferred indication

189	transcription through STAT6 response element in immune cells (such as T- cells).	transcription through the Signal Transducers and Activators of Transcription (STAT6) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT6 transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521- 1523 (2000); Curiel et al., Eur J Immunol 27(8):1982-1987	is allergy. Another highly preferred indication is asthma. Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative
-----	--	--	--

				<p>(1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
190	HMDAB29	604	<p>Regulation of viability and proliferation of pancreatic beta cells.</p>	<p>Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic</p>

			<p>invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ohtani KI, et al., Endocrinology, 139(1):172-8 (1998); Krautheim A, et al, Exp Clin Endocrinol Diabetes, 107 (1):29-34 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that</p>	<p>nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious</p>
--	--	--	---	---

			may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.	Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
190	HMDAB29	604	Insulin Secretion	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel</p>

			<p>disregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell</p>	<p>blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred</p>
--	--	--	--	--

				line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.	indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
190	HMDAB29	604	CD152 in Human T cells		
190	HMDAB29	604	SEAP in SW480		
191	HMDAD44	605	Regulation of transcription via DMEF1 response element in adipocytes and pre-adipocytes	Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate	A highly preferred indication is diabetes mellitus. Additional highly preferred indications include complications associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic

				<p>insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and,</p>	<p>neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively,</p>
--	--	--	--	--	---

				<p>Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.</p>	<p>weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
191	HMDAD44	605	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly</p>

				<p>(including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety.</p>	<p>preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis,</p>
--	--	--	--	--	--

				Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
191	HMDAD44	605	Production of TNF alpha by dendritic cells	TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing)

			<p>cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFα), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol</p>	<p>TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms</p>
--	--	--	--	--

			<p>158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious</p>
--	--	--	---	--

192	HMEBB82	606	Production of MIP1alpha	<p>MIP-1alpha FMT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders.</p>
-----	---------	-----	-------------------------	--	--

				<p>invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy.</p> <p>Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders").</p> <p>Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer.</p> <p>Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	---	---

					hyperplasia, metaplasia, and/or dysplasia.
193	HMEDE24	607	IgG in Human B cells SAC		
193	HMEDE24	607	Activation of transcription through NFAT response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA	Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as

			<p>85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	---	---

194	HMELM75	608	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia,</p>
-----	---------	-----	---------------------	--	---

				<p>204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	--	---	--

195	HMI AK10	609	<p>Activation of transcription through cAMP response element (CRE) in pre-adipocytes.</p>	<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. For example, a 3T3-L1/CRE reporter assay may be used to identify factors that activate the cAMP signaling pathway. CREB plays a major role in adipogenesis, and is involved in differentiation into adipocytes. CRE contains the binding sequence for the transcription factor CREB (CRE binding protein). Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies</p>	<p>dysplasia.</p> <p>A highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. An additional highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease,</p>
-----	----------	-----	---	---	---

			and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Reusch et al., Mol Cell Biol 20(3):1008-1020 (2000); and Klemm et al., J Biol Chem 273:917-923 (1998), the contents of each of which are herein incorporated by reference in its entirety. Pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.	hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). Additional highly preferred indications are complications associated with insulin resistance.
HMI/AK10	609	Activation of	Assays for the activation of	Highly preferred indications

195	transcription through STAT6 response element in immune cells (such as mast cells).	transcription through the Signal Transducers and Activators of Transcription (STAT6) response element in immune cells (such as in the human HMC-1 mast cell line) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT6 transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Sherman, Immunol Rev 179:48-56 (2001); Malaviya and Uekun, J Immunol 168:421-426 (2002);	include allergy, asthma, and rhinitis. Additional highly preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include hematopoietic and immunological disorders (e.g., as described below under "Immune Activity", "Blood- Related Disorders", and/or "Cardiovascular Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancer, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other
-----	--	---	--

				<p>Masuda et al., J Biol Chem 275(38):29331-29337 (2000); and Masuda et al., J Biol Chem 276:26107-26113 (2001), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include hematopoietic and immunological disorders such as arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
195	HMIK10	609	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune</p>

				<p>in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid</p>
--	--	--	--	---	---

					<p>tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	--	---

195	HMI AK10	609	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications</p>
-----	----------	-----	--	--	---

<p>155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>

195	HMI AK10	609	<p>Activation of transcription through NFKB response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al.,</p>	<p>meningitis, Lyme Disease, and asthma and allergy.</p> <p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell</p>
-----	----------	-----	---	---	--

				<p>29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
195	HMIK10	609	<p>Activation of transcription through STAT6 response element in immune</p>	<p>Assays for the activation of transcription through the Signal Transducers and Activators of Transcription (STAT6) response</p>	<p>A highly preferred indication is allergy. Another highly preferred indication is asthma. Additional highly</p>

			<p>cells (such as T-cells).</p>	<p>element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT6 transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521-1523 (2000); Curriel et al., Eur J Immunol 27(8):1982-1987 (1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of</p>	<p>preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	---------------------------------	--	---

				<p>which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
196	HMIBD93	610	IFNg in Human T-cell 2B9		
196	HMIBD93	610	VEGF in SW480		
196	HMIBD93	610	Production of IL-13 and activation of T-cells.	Assays for production of IL-13 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as

				<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-13 and/or activation of T-cells. Exemplary assays for IL-13 production that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays such as disclosed and/or cited in: Grunig, G, et al., "Requirement for IL-13 independently of IL-4 in Experimental asthma" Science;282: 2261-2263 (1998), and Wills-Karp M, et al., "Interleukin-13: central mediator of allergic asthma" Science; 282: 2258-2261 (1998); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL13, a Th2 type cytokine, is a potent stimulus for mucus production, airway hyper-responsiveness</p>	<p>described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	--	--

197	HMIBF07	611	Production of MCP-1	<p>and allergic asthma. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p> <p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood</p>
-----	---------	-----	---------------------	--	---

			<p>surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremim, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or</p>	<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases</p>
--	--	--	--	--

				cytokines, initiate and upregulate T cell proliferation and functional activities.	(e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
197	HMIBF07	611	Insulin Secretion	Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes.	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic

				<p>Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed</p>	<p>neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain.</p>
--	--	--	--	---	---

197	HMIBF07	611	Stimulation of insulin secretion from pancreatic beta cells.	<p>with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin</p>	<p>Additional highly preferred indications are complications associated with insulin resistance.</p>
				<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness,</p>	

				<p>secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al, FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety.</p> <p>Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated.</p> <p>Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References:</p>	<p>nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin</p>
--	--	--	--	--	---

				Asfari et al. Endocrinology 1992 130:167.	resistance.
198	HMICP65	612	HLA-DR in Human T cells	Assays for measuring expression of VCAM are well- known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106),	Highly preferred indications include inflammation (acute and chronic), restnosis, atherosclerosis, asthma and allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood- Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,
198	HMICP65	612	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells (HUVEC))		

198	HMICP65	612	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.</p>	<p>hyperplasia, metaplasia, and/or dysplasia.</p>
			<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications</p>	

				<p>antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia</p>
--	--	--	--	---	--

200.	HMSBE04	614	Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays	(ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
				Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as	

				<p>disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs</p>
--	--	--	--	--	---

201	HMSCL38	615	Activation of transcription through NFKB response element in immune cells (such as B-cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gri G, et al., Biol Chem, 273(11):6431-6438 (1998); Pyatt DW, et al., Cell Biol Toxicol 2000;16(1):41-51 (2000); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al.,	and tissues, endocarditis, meningitis, and Lyme Disease. Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma
-----	---------	-----	--	--	---

202	HMSCR69	616		<p>Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Reh B-cell line.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred</p>
			Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate</p>	

			<p>immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p>	<p>indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis,</p>
--	--	--	---	--

				Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
203	HMSHU20	617	Production of IFNgamma using a T cells	IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of	A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related

			<p>inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin</p>	<p>Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for</p>
--	--	--	--	--

				<p>Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
204	HMSHY25	618	SEAP in HIB/CRE		
	HMSHY25	618	Proliferation of pre-	Assays for the regulation (i.e.	

204			adipose cells (such as 3T3-L1 cells)	<p>increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.</p>	
	HMSHY25	618	Production of GM-	GM-CSF FMAT. GM-CSF is	A highly preferred

204			CSF	<p>expressed by activated T cells, macrophages, endothelial cells, and fibroblasts. GM-CSF regulates differentiation and proliferation of granulocytes-macrophage progenitors and enhances antimicrobial activity in neutrophils, monocytes and macrophage. Additionally, GM-CSF plays an important role in the differentiation of dendritic cells and monocytes, and increases antigen presentation. GM-CSF is considered to be a proinflammatory cytokine. Assays for immunomodulatory proteins that promote the production of GM-CSF are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and modulate the growth and differentiation of leukocytes. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as GM-CSF, and the activation of T cells.</p>	<p>embodiment of the invention includes a method for stimulating the production of GM-CSF. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of GM-CSF. Highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., as described below under "Infectious Disease". Highly preferred indications include blood disorders (e.g., neutropenia (and the prevention of neutropenia (e.g., in HIV infected patients), and/or as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include asthma. Highly</p>
-----	--	--	-----	---	--

			<p>Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Ye et al., J Leukoc Biol (58(2):225-233, the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC) or may be isolated using techniques disclosed herein or otherwise known in the art. Natural killer (NK) cells are large granular lymphocytes that have cytotoxic activity but do bind antigen. NK cells show antibody-independent killing of tumor cells and also recognize antibody bound on target cells, via NK Fc receptors, leading to cell-mediated cytotoxicity.</p>	<p>preferred indications include neoplastic diseases (e.g., leukemia (e.g., acute lymphoblastic leukemia, and acute myelogenous leukemia), lymphoma (e.g., non-Hodgkin's lymphoma and Hodgkin's disease), and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications include: suppression of immune reactions to transplanted organs and tissues (e.g., bone marrow transplant); accelerating myeloid recovery; and mobilizing hematopoietic progenitor cells. Preferred indications include boosting a T cell-mediated immune response, and alternatively, suppressing a T cell-mediated immune</p>
--	--	--	---	---

				<p>response. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and allergy.</p>
204	HMSHY25	618	<p>Production of IFNgamma using Natural Killer cells</p>	<p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2; promotes IgG2a and inhibits IgE; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of</p> <p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" (e.g. cancer/tumorigenesis) and/or "Cardiovascular Disorders"),</p>

				<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology</p>	<p>and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response, boosting antibody-dependent immune responses, suppressing antibody-dependent immune responses, boosting innate immunity and immune responses, and suppressing innate immunity and immune responses. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma,</p>
--	--	--	--	--	--

				<p>(Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Natural Killer (NK) cells that may be used according to these assays are publicly available (e.g., through the ATCC) or may be isolated using techniques disclosed herein or otherwise known in the art. Natural killer (NK) cells are large granular lymphocytes that have cytotoxic activity but do bind antigen. NK cells show antibody-independent killing of tumor cells and also recognize antibody bound on target cells, via NK Fc receptors, leading to cell-mediated cytotoxicity.</p>	<p>melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	---	---

205	HMTAB77	619	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T</p>	<p>asthma and allergy.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients</p>
-----	---------	-----	--	---	--

				<p>cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	---	---

				neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
206	HMUAE26	620	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for</p>

			<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and</p>
--	--	--	---	---

					atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous),
--	--	--	--	--	---

					<p>glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and</p>
--	--	--	--	--	--

					<p>injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include</p>
--	--	--	--	--	---

206	HMUAE26	620	Production of IFNgamma using a T cells	IFNgamma FMA.T. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention	autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
					<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease</p>

				<p>(including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999),</p>	<p>and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include</p>
--	--	--	--	--	--

				<p>the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
207	HMVDU15	621	<p>Activation of transcription through cAMP response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease").</p>

				<p>invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is a suspension</p>	<p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	---	---

				<p>culture of IL-2 dependent T cells that also respond to IL-4.</p>	<p>hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
208	HMWJF53	622	<p>Inhibition of squalene synthetase gene transcription.</p>	<p>Reporter Assay: construct contains regulatory and coding sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the contents of which are herein incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72</p>	

208	HMW/JF53	622		<p>hours. HepG2 is a human hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.</p> <p>IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an</p>
-----	----------	-----	--	--	---

			<p>stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell</p>	<p>infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	--

				receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
209	HNEAK81	623	Production of MIP1alpha	MIP-1alpha FMA T. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include</p>

				<p>differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells</p>	<p>blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	--	--

				that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
210	HNECL22	624	IgG in Human B cells SAC		
210	HNECL22	624	Activation of transcription through NFKB response element in immune cells (such as T-cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis,

				<p>immunomodulatory genes. Exemplary assays for transcription through the NFkB response element that may be used or routinely modified to test NFkB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic</p>
--	--	--	--	---	--

					anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.
					Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as
					Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention
					Activation of T-Cell p38 or JNK Signaling Pathway.
					625
					HNECW49
211					

				<p>(including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues,</p>
--	--	--	--	---	--

212	HNEDH88	626	<p>Activation of transcription through AP1 response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol</p>	<p>endocarditis, meningitis, and Lyme Disease.</p> <p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers,</p>
-----	---------	-----	--	--	---

				<p>29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
213	HNFAC50	627	<p>Regulation of apoptosis of immune cells (such as mast cells).</p>	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and</p>

				<p>invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E -antigen, promoted by T helper cell type 2 cytokines, is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., J Biol Chem, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., J Exp Med, 192(8):1093-1103 (2000); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents</p>	inflammation.
--	--	--	--	---	---------------

				of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary immune cells that may be used according to these assays include mast cells such as the HMC human mast cell line.	
214	HNFHF34	628	Production of MIP1alpha	MIP-1alpha FMA T. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Highly preferred indications include autoimmune diseases</p>

				<p>the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when</p>	<p>(e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications</p>
--	--	--	--	---	--

				activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
215	HNGAM58	629	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described

				<p>Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	--	---	---

					<p>dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
215	HNGAM58	629		IgG in Human B cells SAC	
215	HNGAM58	629		CD152 in Human T cells	
216	HNGBH53	630		Activation of transcription through CD28 response element in immune cells (such as T-	<p>Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to</p> <p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred</p>

			<p>assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T</p>	<p>embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. An additional highly preferred</p>
--	--	--	---	---

				<p>cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>indication includes infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication is infection (e.g., tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or an infectious disease as described below under</p>
--	--	--	--	--	--

					<p>"Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
217	HNGDQ38	631	Production of MCP-1	MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large	<p>A highly preferred embodiment of the invention includes a method for</p>

				<p>variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160</p>	<p>stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic</p>
--	--	--	--	--	---

217			MIP1alpha	<p>immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al.,</p>	<p>embodiment of the invention includes a method for stimulating MIP 1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP 1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute</p>
-----	--	--	-----------	---	--

				<p>"Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
	HNGDX18	632	Regulation of	Assays for the regulation of	A highly preferred indication is

218	transcription via DMEF1 response element in adipocytes and pre-adipocytes	transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M. V., et al., J	diabetes mellitus. Additional highly preferred indications include complications associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic
-----	---	--	--

				<p>Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under</p>	<p>retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	--

218	HNGDX18	632	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988);</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response,</p>
-----	---------	-----	--	---	---

				<p>Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs</p>
--	--	--	--	--	--

218	HNGDX18	632	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black</p>	<p>and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional</p>
-----	---------	-----	--	--	--

				<p>et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas,</p>
--	--	--	--	---	---

					multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
219	HNGDY34	633	Production of IL-6	IL-6 F/MAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins	A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related

				<p>produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical</p>	<p>Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia,</p>
--	--	--	--	--	--

				<p>approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
	HNGEA34	634	Production of IL-5	IL-5 FMAT. Assays for	A highly preferred

220				<p>immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>
-----	--	--	--	---	---

				<p>invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>"Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma,</p>
--	--	--	--	--	---

				<p>arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
221	HNGGA68	635	<p>Activation of transcription through cAMP response element (CRE) in pre-adipocytes.</p>	<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. For example, a 3T3-L1/CRE reporter assay may be used to identify factors that activate the cAMP signaling pathway. CREB plays a major role in adipogenesis, and is involved in differentiation into adipocytes. CRE contains the binding sequence for the transcription</p>
				<p>A highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. An additional highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic</p>

				<p>factor CREB (CRE binding protein). Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Reusch et al., Mol Cell Biol 20(3):1008-1020 (2000); and Klemm et al., J Biol Chem 273:917-923 (1998), the contents of each of which are herein incorporated by reference in its entirety. Pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3</p>	<p>neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hypoglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	--

221	HNGGA68	635	<p>Activation of transcription through serum response element in pre-adipocytes.</p>	<p>fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA</p>	<p>A highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. An additional highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness,</p>
-----	---------	-----	--	--	--

				<p>85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. Pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below). Additional highly preferred indications are complications associated with insulin resistance.</p>
221	HNGGA68	635	<p>Activation of transcription through cAMP response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below</p>

				<p>agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the</p>	<p>under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	--	--

				CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.	conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
221	HNGGA68	635	Activation of serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune

				<p>in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid</p>
--	--	--	--	--	---

					<p>tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	--	---

221	HNGGA68	635	<p>Activation of transcription through NFKB response element in immune cells (such as EOL1 cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., <i>Gene</i> 66:1-10 (1998); Cullen and Malm, <i>Methods in Enzymol</i> 216:362-368 (1992); Henthorn et al., <i>Proc Natl Acad Sci USA</i> 85:6342-6346 (1988); Valle Blazquez et al, <i>Immunology</i> 90(3):455-460 (1997); Aramburau et al., <i>J Exp Med</i> 82(3):801-810 (1995); and Fraser et al., 29(3):838-844</p>	<p>Highly preferred indications include asthma, allergy, hypersensitivity reactions, and inflammation. Preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), immunological disorders, inflammation and inflammatory disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below).</p>
-----	---------	-----	--	--	---

				<p>(1999), the contents of each of which are herein incorporated by reference in its entirety. For example, a reporter assay (which measures increases in transcription inducible from a NFkB responsive element in EOL-1 cells) may link the NFkB element to a reporter gene and binds to the NFkB transcription factor, which is upregulated by cytokines and other factors. Exemplary immune cells that may be used according to these assays include eosinophils such as the human EOL-1 cell line of eosinophils. Eosinophils are a type of immune cell important in the allergic responses; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Eol-1 is a human eosinophil cell line.</p>	
221	HNGGA68	635	<p>Activation of transcription through GATA-3 response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the GATA-3 signaling pathway in HMC-1 human mast cell line. Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and</p>

			<p>for the activation of transcription through the GATA3 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J</p>	<p>inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia</p>
--	--	--	---	---

				<p>Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and Henderson et al., Mol Cell Biol 14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>(ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
221	HNGGA68	635	<p>Activation of transcription through NFAT response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the NFAT signaling pathway in HMC-1 human mast cell line. Activation of NFAT in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under</p>

				<p>are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-7223 (2000); Hutchinson and McCloskey, J Biol Chem 270(27):16333-16338 (1995), and Turner et al., J Exp Med 188:527-537 (1998), the</p>	<p>"Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel</p>
--	--	--	--	--	--

				contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.	disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
221	HNGGA68	635	MCP-1 in HUVEC		
221	HNGGA68	635	SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)		
221	HNGGA68	635	Activation of transcription through NFKB response element in immune cells (such as basophils).	This reporter assay measures activation of the NFKB signaling pathway in Ku812 human basophil cell line. Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and	Highly preferred indication includes allergy, asthma, and rhinitis. Additional highly preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include immunological and hematopoietic disorders (e.g.,

				<p>agonists or antagonists of the invention) to regulate NFκB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFκB response element that may be used or routinely modified to test NFκB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Marone et al, Int Arch Allergy Immunol 114(3):207-17 (1997), the contents of each of which are herein incorporated by reference in its entirety. Basophils that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human basophil cell lines that may be used according to these assays include Ku812, originally established from a patient with chronic</p>	<p>as described below under "Immune Activity", and "Blood-Related Disorders"). Preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancer, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary tract cancers and as described below under "Hyperproliferative Disorders".</p>
--	--	--	--	---	---

				myelogenous leukemia. It is an immature prebasophilic cell line that can be induced to differentiate into mature basophils.	
221	HNGGA68	635	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al.,	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional

				<p>Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas,</p>
--	--	--	--	--	---

				multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
222	HNGIV64	636	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity</p> <p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method</p>

				<p>that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-</p>	<p>for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural</p>
--	--	--	--	--	---

				adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.	disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as
--	--	--	--	---	---

					described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia,
--	--	--	--	--	---

					gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
222	HNGIV64	636	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as

				<p>used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated</p>	<p>described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	---	--

				via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
223	HNGJB41	637	Production of ICAM-1	<p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p>	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke

224	HNGKT4I	638	SEAP in 293/ISRE		
224	HNGKT4I	638	Activation of transcription through cAMP response element (CRE) in pre-adipocytes.	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. For example, a 3T3-L1/CRE reporter assay may be used to identify factors that activate the cAMP signaling pathway. CREB plays a major role in adipogenesis, and is involved in differentiation into adipocytes. CRE contains the binding sequence for the transcription factor CREB (CRE binding protein). Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the	A highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. An additional highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis,

				<p>invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Reusch et al., Mol Cell Biol 20(3):1008-1020 (2000); and Klemm et al., J Biol Chem 273:917-923 (1998), the contents of each of which are herein incorporated by reference in its entirety. Pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---	---

224	HNGKT41	638	<p>Activation of transcription through NFKB response element in immune cells (such as basophils).</p>	<p>This reporter assay measures activation of the NFKB signaling pathway in Ku812 human basophil cell line. Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Marone et al, Int Arch Allergy Immunol 114(3):207-17 (1997), the</p>	<p>Highly preferred indication includes allergy, asthma, and rhinitis. Additional highly preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include immunological and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"). Preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancer, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic,</p>
-----	---------	-----	---	---	--

				contents of each of which are herein incorporated by reference in its entirety. Basophils that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human basophil cell lines that may be used according to these assays include Ku812, originally established from a patient with chronic myelogenous leukemia. It is an immature prebasophilic cell line that can be induced to differentiate into mature basophils.	esophageal, stomach, brain, liver, urinary tract cancers and as described below under "Hyperproliferative Disorders".
224	HNGKT41	638	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications

				<p>the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other</p>
--	--	--	--	--	---

					<p>preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
224	HNGKT41	638	Activation of transcription through API response element in immune	Assays for the activation of transcription through the API response element are well-known in the art and may be	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"),</p>

			<p>used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly</p>	<p>blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	---

				<p>available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
224	HNGKT41	638	<p>Activation of transcription through CD28 response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells.</p>

			<p>test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g.,</p>
--	--	--	---	---

					metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also
--	--	--	--	--	--

224	HNGKT41	638	Activation of transcription through NFAT response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to	include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
				Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred	

				<p>test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease,</p>
--	--	--	--	--	--

224	HNGKT41	638	Activation of transcription through NFKB response element in immune cells (such as T-cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays	inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
				Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred	

				<p>disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	--	--

225	HNGNK44	639	Production of TNF alpha by dendritic cells	<p>TNFa F/MAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and</p>	<p>suppression of immune reactions to transplanted organs, asthma and allergy.</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and</p>
-----	---------	-----	--	--	---

			<p>agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease,</p>
--	--	--	---	--

225	HNGNK44	639	Proliferation of pre-adipose cells (such as 3T3-L1 cells)	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on	inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
-----	---------	-----	---	---	---

				quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.	
225	HNGNK44	639	IgG in Human B cells		
225	HNGNK44	639	CD152 in Human T cells		
226	HNGNO53	640	Inhibition of squalene synthetase gene transcription.	Reporter Assay: construct contains regulatory and coding sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the contents of which are herein incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72 hours. HepG2 is a human	

226	HNGNO53	640	Activation of transcription through serum response element in immune cells (such as T-cells).	hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated</p>
-----	---------	-----	---	---	--

			<p>368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety.</p> <p>Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders").</p> <p>Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas,</p>
--	--	--	---	---

					multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
226	HNGN053	640	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T

				<p>the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes</p>	<p>cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	--	--

227	HNGPJ25	641	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>isolated from cord blood.</p> <p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders").</p> <p>Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-</p>
-----	---------	-----	--	---	--

				<p>93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia,</p>
--	--	--	--	---	--

227	HNGPJ25	641	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-	hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy. A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications
-----	---------	-----	---	--	---

			<p>117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's</p>
--	--	--	--	--

					lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
227	HNGPJ25	641	Regulation of transcription of Malic Enzyme in adipocytes	Assays for the regulation of transcription of Malic Enzyme are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate transcription of Malic Enzyme, a key enzyme in lipogenesis. Malic enzyme is involved in lipogenesis and its expression is stimulated by insulin. ME promoter contains two direct repeat (DR1)-like elements	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke,</p>

				<p>MEp and MEd identified as putative PPAR response elements. ME promoter may also responds to API and other transcription factors.</p> <p>Exemplary assays that may be used or routinely modified to test for regulation of transcription of Malic Enzyme (in adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Streeter, R.S., et al., Mol Endocrinol, 12(11):1778-91 (1998); Garcia-Jimenez, C., et al., Mol Endocrinol, 8(10):1361-9 (1994); Barroso, I., et al., J Biol Chem, 274(25):17997-8004 (1999); Ijpenberg, A., et al., J Biol Chem, 272(32):20108-20117 (1997); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated.</p>	<p>impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss</p>
--	--	--	--	--	---

228	HNHCT47	642	Regulation of viability or proliferation of immune cells (such as human eosinophil EOL-1 cells).	Exemplary hepatocytes that may be used according to these assays includes the H4IIE rat liver hepatoma cell line.	or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Highly preferred indications include eosinophilia, asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications
-----	---------	-----	--	---	--

				<p>Eosinophil cell lines that may be used according to these assays are publicly available and/or may be routinely generated. Exemplary eosinophil cells that may be used according to these assays include EOL-1 Cells.</p>	<p>include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>
228	HNHCT47	642	<p>Production of IL-10 and activation of T-cells.</p>	<p>Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>

				<p>helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p>	
229	HNHFE71	643	<p>Activation of Adipocyte PI3 Kinase Signalling Pathway</p>	<p>Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of</p>	<p>A highly preferred embodiment of the invention includes a method for increasing adipocyte survival An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival.</p>

				<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal</p>	<p>A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related</p>
--	--	--	--	--	--

				isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.	Disorders”), immune disorders (e.g., as described below under “Immune Activity”), neural disorders (e.g., as described below under “Neural Activity and Neurological Diseases”), and infection (e.g., as described below under “Infectious Disease”). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the “Renal Disorders” section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis,
--	--	--	--	---	---

					<p>microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies,</p>
--	--	--	--	--	--

					<p>muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer.</p> <p>Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
229	HNHFE71	643	<p>Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells</p>	<p>Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention</p>	<p>Highly preferred indications include inflammation (acute and chronic), restenosis, atherosclerosis, asthma and allergy. Highly preferred indications include</p>

			(HUVVEC))	<p>(including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.</p>	<p>inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	-----------	--	---

229	HNHFE71	643	Calcium flux in immune cells (such as monocytes)	<p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux in immune cells (such as monocytes) include assays disclosed in: Chan, CC, et al., J Pharmacol Exp Ther, 269(3):891-896 (1994); Andersson, K, et al., Cytokine, 12(12):1784-1787 (2000); Scully, SP, et al., J Clin Invest, 74(2) 589-599 (1984); and, Sullivan, E, et al., Methods Mol Biol, 114:125-133 (1999), the contents of each of which is herein incorporated by reference</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Infection, Inflammation, Atherosclerosis, Hypersensitivity, and Leukemias</p>
-----	---------	-----	--	--	--

				in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the THP-1 monocyte cell line.	
230	HNHGK22	644	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm,	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below),

				<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia,</p>
--	--	--	--	--	---

					<p>pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
231	HNHHB10	645	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins</p>

			<p>transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOL T4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with</p>
--	--	--	--	--

					chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
232	HNHK174	646	Glucose Production in H4IIE		
232	HNHK174	646	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells)	Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention	Highly preferred indications include inflammation (acute and chronic), restenosis, atherosclerosis, asthma and allergy. Highly preferred indications include

			<p>(including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.</p>	<p>inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	---	---

232	HNHKJ74	646	<p>Production of IL-10 and activation of T-cells.</p>	<p>Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
-----	---------	-----	---	--	--

				these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
233	HNTBT17	647	IL-10 in Human T cells	Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases
234	HNTMH79	648	Activation of transcription through API response element in immune cells (such as T-cells).		

				<p>test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the HT2 cell line, which is an IL-2 dependent suspension culture cell line that also responds to IL-4.</p>	<p>(e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's</p>
--	--	--	--	---	--

					lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
235	HODAG07	649	SEAP in ATP-3T3-L1		
236	HODBB70	650	Activation of transcription through cAMP response element in immune cells (such as T-cells).	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred

				<p>disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs</p>
--	--	--	--	---	---

				and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
236	HODBB70	650	Regulation of transcription through the PEPCK promoter in hepatocytes	<p>Assays for the regulation of transcription through the PEPCK promoter are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the PEPCK promoter in a reporter construct and regulate liver gluconeogenesis. Exemplary assays for regulation of transcription through the PEPCK promoter that may be used or routinely modified to test for PEPCK promoter activity (in hepatocytes) of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as</p>

			<p>et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Lochhead et al., Diabetes 49(6):896-903 (2000); and Yeagley et al., J Biol Chem 275(23):17814-17820 (2000), the contents of each of which is herein incorporated by reference in its entirety. Hepatocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary liver hepatoma cells that may be used according to these assays include H4Ile cells, which contain a tyrosine amino transferase that is inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., an infectious diseases or disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred</p>
--	--	--	--	---

					<p>indications include glycogen storage disease (e.g., glycogenoses), hepatitis, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism, and hepatocarcinomas.</p> <p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), infection (e.g., an infectious disease and/or disorder as described below under "Infectious Disease"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases").</p> <p>Additional preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders").</p>
--	--	--	--	--	--

237	HODBV05	651	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell</p>	<p>Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, and urinary cancer. A highly preferred indication is liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood</p>
-----	---------	-----	---------------------	---	--

			<p>surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or</p>	<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases</p>
--	--	--	--	--

				cytokines, initiate and upregulate T cell proliferation and functional activities.	(e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
238	HODCZ32	652	SEAP in HepG2/Squalen synthetase(stimulation)		
238	HODCZ32	652	Activation of Natural Killer Cell ERK Signaling Pathway.	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and	A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative

			<p>differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications</p>
--	--	--	---	--

					include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.
239	HOFNU55	653	Activation of transcription through cAMP response element in immune cells (such as T-	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or

				<p>assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, bind to CREB transcription factor, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the</p>	<p>"Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other</p>
--	--	--	--	---	---

				ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.	preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
239	HOFNU55	653	SEAP in OE-33		
240	HOGBF01	654	Activation of transcription through AP1 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular

				<p>and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., <i>Gene</i> 66:1-10 (1988); Cullen and Malm, <i>Methods in Enzymol</i> 216:362-368 (1992); Henthorn et al., <i>Proc Natl Acad Sci USA</i> 85:6342-6346 (1988); Rellahan et al., <i>J Biol Chem</i> 272(49):30806-30811 (1997); Chang et al., <i>Mol Cell Biol</i> 18(9):4986-4993 (1998); and Fraser et al., <i>Eur J Immunol</i> 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell</p>	<p>Disorders”, and/or “Blood-Related Disorders”), and infection (e.g., an infectious disease as described below under “Infectious Disease”). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred</p>
--	--	--	--	--	---

				line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
240	HOGBF01	654	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by	A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described

				<p>cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925</p>	<p>below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach,</p>
--	--	--	--	--	--

				<p>(1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
240	HOGBF01	654	Production of IFNgamma using a T cells	IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine.	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of</p>

				<p>IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNγ), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention</p>	<p>IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic</p>
--	--	--	--	---	--

			<p>(including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs</p>
--	--	--	--	---

241	HORBS82	655	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999);</p>	<p>and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly</p>
-----	---------	-----	---	--	--

				<p>Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity", and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic</p>
--	--	--	--	--	--

					<p>nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious</p>
--	--	--	--	--	---

					<p>Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and</p>
--	--	--	--	--	---

241	HORBS82	655	Activation of JNK Signaling Pathway in immune cells (such as eosinophils).	Kinase assay. JNK kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be used or routinely modified to test JNK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999);	urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
				Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under	

				<p>Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate signal transduction, cell proliferation, activation, or apoptosis in eosinophils include assays disclosed and/or cited in: Zhang JP, et al., "Role of caspases in dexamethasone-induced apoptosis and activation of c-Jun NH2-terminal kinase and p38 mitogen-activated</p>	<p>"Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>
--	--	--	--	---	---

				<p>protein kinase in human eosinophils" Clin Exp Immunol; Oct;122(1):20-7 (2000); Hebestreit H, et al., "Disruption of fas receptor signaling by nitric oxide in eosinophils" J Exp Med; Feb 2;187(3):415-25 (1998); J Allergy Clin Immunol 1999 Sep;104(3 Pt 1):565-74; and, Sousa AR, et al., "In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-terminal kinase phosphorylation" J Allergy Clin Immunol; Sep;104(3 Pt 1):565-74 (1999); the contents of each of which are herein incorporated by reference in its entirety.</p>	
241	HORBS82	655	<p>Regulation of apoptosis of immune cells (such as mast cells).</p>	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and inflammation.</p>

				<p>example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E -antigen, promoted by T helper cell type 2 cytokines, is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., J Biol Chem, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., J Exp Med, 192(8):1093-1103 (2000); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may</p>
--	--	--	--	---

241	HORBS82	655	<p>Production of IL-13 and activation of T-cells.</p>	<p>be used according to these assays are publicly available (e.g., through commercial sources). Exemplary immune cells that may be used according to these assays include mast cells such as the HMC human mast cell line.</p> <p>Assays for production of IL-13 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-13 and/or activation of T-cells.</p> <p>Exemplary assays for IL-13 production that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays such as disclosed and/or cited in: Grunig, G, et al., "Requirement for IL-13 independently of IL-4 in Experimental asthma"</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
-----	---------	-----	---	---	--

				Science;282: 2261-2263 (1998), and Wills-Karp M, et al., "Interleukin-13: central mediator of allergic asthma" Science; 282: 2258-2261 (1998); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL13, a Th2 type cytokine, is a potent stimulus for mucus production, airway hyper-responsiveness and allergic asthma. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection,
242	HORBV76	656	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of	

				polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
242	HORBV76	656	VEGF in SW480		
243	HOSEC25	657	IL-2 in Human T cells		
244	HOSEI81	658	Activation of T-Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as

				<p>art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the</p>	<p>described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example,</p>
--	--	--	--	--	---

245	HOSE194	659		<p>ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
			Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>

				<p>cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160</p>	<p>"Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and</p>
--	--	--	--	---	---

				<p>(2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
245	HOSEJ94	659	IFNg in Human T-cell 2B9	

246	HOUCA21	660	Production of IFNgamma using a T cells	<p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred</p>
-----	---------	-----	--	--	--

				<p>assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated</p>	<p>indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	---	---

				immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
247	HOUDE92	661	SEAP in 293/ISRE		
247	HOUDE92	661	Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications

			<p>Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
--	--	--	--	--

247	HOUDE92	661	<p>Activation of transcription through cAMP response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-</p>
-----	---------	-----	---	---	--

				<p>which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
247	HOUE92	661	SEAP in HIB/CRE		
247	HOUE92	661	Activation of transcription through	This reporter assay measures activation of the GATA-3	Highly preferred indications include allergy, asthma, and

			<p>GATA-3 response element in immune cells (such as mast cells).</p>	<p>signaling pathway in HMC-1 human mast cell line. Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the GATA3 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn</p>	<p>rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or</p>
--	--	--	--	--	---

				<p>et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and Henderson et al., Mol Cell Biol 14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
247	HOUDE92	661	<p>Activation of transcription through NFAT response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the NFAT signaling pathway in HMC-1 human mast cell line. Activation of NFAT in mast cells has been linked to cytokine and</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under</p>

				chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-		chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-	"Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease,
--	--	--	--	---	--	---	--

				<p>7223 (2000); Hutchinson and McCloskey, J Biol Chem 270(27):16333-16338 (1995), and Turner et al., J Exp Med 188:527-537 (1998), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
247	HOUDE92	661	<p>Activation of transcription through NFAT response in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as</p>

				<p>transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T</p>	<p>described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia,</p>
--	--	--	--	--	--

				cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.	pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
247	HOUDE92	661	SEAP in Jurkat/IL4 promoter		
247	HOUDE92	661	SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)		
247	HOUDE92	661	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include

			<p>response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and</p>	<p>blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma,</p>
--	--	--	--	--

					<p>cytotoxic activity.</p>	<p>melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below</p>
--	--	--	--	--	----------------------------	---

247	HOUDE92	661	<p>Activation of transcription through NFKB response element in immune cells (such as B-cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gri G, et al., Biol Chem, 273(11):6431-6438 (1998); Pyatt DW, et al., Cell Biol Toxicol 2000;16(1):41-51 (2000); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA</p>	<p>under "Infectious Disease"). Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma</p>
-----	---------	-----	---	--	--

247	HOUDE92	661	<p>Activation of transcription through NFKB response element in neuronal cells (such as SKNMC cells).</p>	<p>85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary immune cells that may be used according to these assays include the Reh B-cell line.</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Neurological Diseases and Disorders (e.g. Alzheimer's Disease, Parkinson's Disease, Brain Cancer, Seizures).</p>
-----	---------	-----	---	---	---

				test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Gill JS, et al., Neurobiol Dis, 7(4):448-461 (2000); Tamatani M, et al., J Biol Chem, 274(13):8531-8538 (1999); Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Neuronal cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary neuronal cells that may be used according to these assays include the SKNMC neuronal cell line.	Assays for the activation of	Preferred indications include
HOUDE92	661	Activation of				

247	transcription through API response element in immune cells (such as T-cells).	transcription through the API response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference	neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer.
-----	---	--	--

				<p>in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
247	HOUDE92	661	<p>Activation of transcription through CD28 response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of</p>

				<p>transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers,</p>
--	--	--	--	---	---

					<p>such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related</p>
--	--	--	--	--	--

247	HOUDE92	661	<p>Activation of transcription through NFAT response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for</p>	<p>Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p> <p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response,</p>
-----	---------	-----	---	---	---

			transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and	and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas,
--	--	--	--	--

				IL-4 responsive T cells.	multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
247	HOUDE92	661	Activation of transcription through NFKB response element in immune cells (such as T-cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an

				<p>invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis,</p>
--	--	--	--	---	---

247	HOUDE92	661	<p>Activation of transcription through STAT6 response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Signal Transducers and Activators of Transcription (STAT6) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT6 transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol</p>	<p>hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p> <p>A highly preferred indication is allergy. Another highly preferred indication is asthma. Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred</p>
-----	---------	-----	--	---	--

			<p>216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521-1523 (2000); Curiel et al., Eur J Immunol 27(8):1982-1987 (1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious</p>
--	--	--	---	--

248	HOUED72	662	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the</p>	<p>Disease"). Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma,</p>
-----	---------	-----	--	--	--

				contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
248	HOUED72	662	SEAP in UMR-106		
249	HOUFS04	663	CD152 in Human T cells		
249	HOUFS04	663	Activation of transcription through GAS response element in immune cells (such as T-	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders").

			cells).	<p>be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the</p>	<p>Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune</p>
--	--	--	---------	--	--

				<p>MOLT4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
249	HOUFS04	663	Activation of transcription through	Assays for the activation of transcription through the NFkB	Highly preferred indications include inflammation and

			<p>NFKB response element in immune cells (such as T-cells).</p>	<p>response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the</p>	<p>inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other</p>
--	--	--	---	--	--

				MOLT4, that may be used according to these assays are publicly available (e.g., through the ATCC).	preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.
250	HOUHI25	664	Activation of transcription through STAT6 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Signal Transducers and Activators of Transcription (STAT6) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and	A highly preferred indication is allergy. Another highly preferred indication is asthma. Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described

				<p>agonists or antagonists of the invention) to regulate STAT6 transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521-1523 (2000); Curiel et al., Eur J Immunol 27(8):1982-1987 (1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the</p>	<p>below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute</p>
--	--	--	--	---	--

				ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.	lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
251	HOVBD85	665	IFNg in Human T-cell 2B9		
251	HOVBD85	665	IL-2 in Human T-cell 2B9		
251	HOVBD85	665	TNFa in Human T-cell 2B9		
251	HOVBD85	665	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing)

				<p>agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell</p>	<p>TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms</p>
--	--	--	--	---	---

				<p>line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is</p>
--	--	--	--	---	--

					infection (e.g., an infectious disease as described below under "Infectious Disease").
251	HOVBD85	665	ICAM in Normal Human Bronchial Epitheliae		
252	HPCAL26	666	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm,	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis

				<p>Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	--	--

				<p>sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
252	HPCAL26	666	<p>Production of IL-10 and activation of T-cells.</p>	<p>Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br</p>
				<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>

				<p>Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p>	
253	HPEBA84	667	<p>Production of IL-10 and activation of T-cells.</p>	<p>Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention</p>	<p>Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune</p>

				<p>(including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6.</p>	<p>Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	---	--

254	HPFBA54	668	<p>Activation of transcription through AP1 response element in immune cells (such as T-cells).</p>	<p>Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and</p>
-----	---------	-----	--	---	--

				<p>216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
--	--	--	--	---	---

254	HPFBA54	668	Production of IFNgamma using a T cells	<p>IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNγ), and the activation of T cells. Such</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred</p>
-----	---------	-----	--	--	--

				assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated	indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,
--	--	--	--	--	--

				immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
254	HPFBA54	668	IL-8 in Normal Human Bronchial Epitheliae		
255	HPFCI36	669	Production of MIP1alpha	MIP-1alpha FMA T. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory	A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications

			<p>protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders").</p>
--	--	--	---	---

				<p>suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
255	HPFC136	669	SEAP in HIB/CRE		
255	HPFC136	669	Stimulation of Calcium Flux in pancreatic beta cells.	<p>Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel</p>

				<p>and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 (Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and</p>	<p>blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred</p>
--	--	--	--	---	--

				glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.	indications are complications associated with insulin resistance.
255	HPFC136	669	Activation of Skeletal Muscle Cell PI3 Kinase Signalling Pathway	Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol	A highly preferred embodiment of the invention includes a method for increasing muscle cell survival An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention

				<p>Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	<p>includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is inhibited. Highly preferred indications include disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under</p>
--	--	--	--	---	---

					<p>"Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g. due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section</p>
--	--	--	--	--	--

					<p>below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity.</p> <p>Additional highly preferred indications include weight loss or alternatively, weight gain.</p> <p>Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal system including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities,</p>
--	--	--	--	--	---

					heart disease, cardiac arrest, heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.
256	HPJBU43	670	CD152 in Human T cells		
256	HPJBU43	670	CD69 in Human T cells		
257	HPMBX22	671	Regulation of viability or proliferation of immune cells (such as human eosinophil EOL-1 cells).	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of eosinophil cells and cell lines.	Highly preferred indications include eosinophilia, asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid

				<p>For example, the CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Eosinophils are a type of immune cell important in allergic responses; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Eosinophil cell lines that may be used according to these assays are publicly available and/or may be routinely generated. Exemplary eosinophil cells that may be used according to these assays include EOL-1 Cells.</p>	<p>arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>
257	HPMBX22	671	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease,</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement</p>

				<p>plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described</p>
--	--	--	--	--	---

				<p>invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an</p>
--	--	--	--	--	--

258	HPMCJ84	672	Activation of transcription through CD28 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-	infectious disease as described below under "Infectious Disease"). A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid
-----	---------	-----	--	--	--

				<p>2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. An additional highly preferred indication includes infection (e.g., AIDS, and/or as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	---	--

					<p>conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication is infection (e.g., tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia,</p>
--	--	--	--	--	--

259	HPMCV30	673	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular	neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy. Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
-----	---------	-----	----------------------	--	---

260	HPMIFH77	674	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>endothelial cells (MVEC).</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients</p>
-----	----------	-----	--	--	---

				<p>cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	---	---

261	HPQCC53	675	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell</p>	<p>neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood</p>
-----	---------	-----	---------------------	---	--

			<p>surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or</p>	<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases</p>
--	--	--	--	--

				cytokines, initiate and upregulate T cell proliferation and functional activities.	(e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
262	HPTRM02	676	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include

				<p>may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is</p>
--	--	--	--	---	--

					<p>idiopathic pulmonary fibrosis.</p> <p>Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
263	HPWBA29	677	<p>Activation of transcription through NFAT response in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T</p>

				<p>functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a</p>	<p>cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic</p>
--	--	--	--	--	---

				<p>suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
263	HPWBA29	677	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative</p>

				<p>modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOL T4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis.</p>
--	--	--	--	---	---

				<p>Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
265	HRADA42	679	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or</p>
				<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"),</p>

				<p>routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain,</p>
--	--	--	--	--	--

					<p>liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
265	HRADA42	679	SEAP in HIB/CRE		
	HRADA42	679	IFNg in Human T-		

265	HRADF49	680	cell 2B9 Activation of Adipocyte ERK Signaling Pathway	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and	A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").
266					

				<p>Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the</p>
--	--	--	--	---	--

					<p>"Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is</p>
--	--	--	--	--	--

					<p>obesity and/or complications associated with obesity.</p> <p>Additional highly preferred indications include weight loss or alternatively, weight gain.</p> <p>Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders.</p> <p>Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer.</p> <p>Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign</p>
--	--	--	--	--	--

					dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
266	HRADF49	680	SEAP in Alk Phos C2C12		
266	HRADF49	680	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated

				<p>immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin’s disease, acute lymphocytic</p>
85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.				

					anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
266	HRADF49	680	MIP-1a in HMC		
266	HRADF49	680	IL-2 in Human T-cell 2B9		
267	HRADN25	681	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include

				<p>response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma,</p>
--	--	--	--	---	--

					<p>melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below</p>
--	--	--	--	--	---

268	HRADT25	682	Production of IFNgamma using a T cells	<p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon</p>	<p>under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T</p>
-----	---------	-----	--	--	---

			<p>gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or</p>	<p>cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
--	--	--	---	---

				CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
269	HRDA117	683	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke

				these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	
269	HRDA117	683	SEAP in OE-33		
270	HRDDQ39	684	Regulation of transcription via DMEF1 response element in adipocytes and pre-adipocytes	Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive	A highly preferred indication is diabetes mellitus. Additional highly preferred indications include complications associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,

				<p>glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be</p>	<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	--

					routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.	
270	HRDDQ39	684	Glucose Production in H4IIE			
270	HRDDQ39	684	Stimulation of insulin secretion from pancreatic beta cells.		Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes.	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic

				<p>Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics</p>	<p>neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, <u>weight gain</u>.</p>
--	--	--	--	--	--

				typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Afari et al. Endocrinology 1992 130:167.	Additional highly preferred indications are complications associated with insulin resistance.
270	HRDDQ39	684	SEAP in UMR-106		
271	HRDER22	685	Glucose Production in H4IIE		
272	HRDFK37	686	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below),

				<p>Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia,</p>
--	--	--	--	---	---

					<p>pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
272	HRDFK37	686	Hexosaminidase in RBL-2H3	<p>IFNgamma FMAT. IFNg plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNg promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNg. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNg. Highly preferred</p>
273	HRGBD54	687	Production of IFNgamma using a T cells		

			<p>expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-</p>	<p>indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described</p>
--	--	--	--	--

			<p>204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---

273	HRGBD54	687	SEAP in UMR-106	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and
274	HSAVA08	688	Activation of transcription through serum response element in immune cells (such as T-cells).		

				<p>by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease,</p>
--	--	--	--	---	---

					inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
274	HSAVA08	688	Production of IL-5	IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5 production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production.</p>

			<p>immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used</p>	<p>A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and</p>
--	--	--	---	---

				<p>according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
274	HSAVA08	688	SEAP in Senescence Assay		
275	HSAWN53	689	Production of ICAM-1	Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection,

				<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Rolfe BE, et al., Atherosclerosis, 149(1):99-110 (2000); Panettieri RA Jr, et al., J Immunol, 154(5):2358-2365 (1995); and, Grunstein MM, et al., Am J Physiol Lung Cell Mol Physiol, 278(6):L1154-L1163 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include Aortic Smooth Muscle Cells (AOSMC); such as bovine AOSMC.</p>	<p>diagnosis, prevention, and/or treatment of Vascular Disease, Atherosclerosis, Restenosis, Stroke, and Asthma.</p>
276	HSA WZ40	690	Production of ICAM-1	<p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection,</p>

				polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
277	HSDZM54	691	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production.

			<p>invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural</p>	<p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for</p>
--	--	--	---	---

				<p>killer cell line with cytolytic and cytotoxic activity.</p>	<p>example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious</p>
--	--	--	--	--	---

278	HSHAX04	692	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein	disease as described below under "Infectious Disease"). Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.
-----	---------	-----	--	--	---

				incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
279	HSHBF76	693	Endothelial Cell Apoptosis	Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative

				<p>associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and</p>	<p>highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g.,</p>
--	--	--	--	--	--

				immune cell extravasation.	heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias,
--	--	--	--	----------------------------	--

				<p>and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis,</p>
--	--	--	--	---

					<p>and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood</p>
--	--	--	--	--	---

					<p>disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
279	HSHBF76	693	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as</p>

				<p>immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-</p>	<p>described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms</p>
--	--	--	--	---	---

			<p>204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	---	---

280	HSKDR27	694	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment</p>
-----	---------	-----	----------------------------	---	---

				<p>assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as</p>
--	--	--	--	---	---

					<p>diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative</p>
--	--	--	--	--	---

				<p>disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and</p>
--	--	--	--	---

					<p>coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
--	--	--	--	--	---

280	HSKDR27	694	<p>Activation of T-Cell p38 or JNK Signaling Pathway.</p>	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach,</p>
-----	---------	-----	---	--	---

				are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
280	HSKDR27	694	Production of IL-4	IL-4 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells that stimulate B cells, T cells, macrophages and mast cells and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-4 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-4 production. A highly preferred indication includes asthma. A highly preferred</p>

				<p>(including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cells, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-4, and the stimulation of immune cells, such as B cells, T cells, macrophages and mast cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):277-283 (1994); Yssel et al., Res Immunol 144(8):610-616 (1993); Bagley et al., Nat Immunol 1(3):257-261 (2000); and van der Graaff et al., Rheumatology (Oxford)</p>	<p>indication includes allergy. A highly preferred indication includes rhinitis. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g.,</p>
--	--	--	--	--	---